

HLA-DRB1 allele distribution among children with rheumatic heart disease in Haji Adam Malik Hospital Medan, Indonesia

Tina Christina Lumban Tobing¹, Teddy Ontoseno², Sri Endah Rahayuningsih³, Ratna Akbari Ganie⁴, Yahwardiah Siregar⁵

¹Department of Child Health, School of Medicine, Universitas Sumatera Utara, ²Department of Child Health, School of Medicine, Universitas Airlangga, ³Department of Child Health, School of Medicine, Universitas Padjajaran, ⁴Department of Clinical Pathology, ⁵Department of Biochemistry, School of Medicine, Universitas Sumatera Utara, Indonesia

ABSTRACT

Rheumatic heart disease (RHD) is influenced by genetic factor, microorganism's virulence, and environmental factor. The aim of this study was to determine human leukocyte (HLA)-DRB1 allele among children with RHD in Medan, Indonesia.

Methods An observational study was conducted at the Department of Child Health, Haji Adam Malik Hospital Medan from April to June 2017. Inclusion criteria were children aged 5-18 years diagnosed with RHD. Children with concomitant congenital heart disease were excluded. HLA-DRB1 alleles were analyzed using the PCR sequence-specific priming (SSP) technique. Statistical analysis was done using computer software.

Results A total of 62 children were enrolled. The mean age of children was 12.6 (SD 3.44) years; 33 (53.2%) were male. The most dominant allele was HLA-DRB1*12, followed by HLA-DRB1*15.

Conclusion It is proven in this research that RHD is influenced by genetic factor with HLA-DRB1*12 allele found to be the most common allele in children with RHD in Medan, Indonesia.

Key words: cardiology, human leukocyte antigen, paediatric, RHD

Corresponding author:

Tina Christina Lumban Tobing
Department of Child Health,
School of Medicine,
Universitas Sumatera Utara
17 Bunga Lau Street, Medan 20136,
North Sumatera, Indonesia
Phone: +62 811 653 616;
E-mail: tinatobing61@gmail.com

Original submission:

18 November 2019;

Revised submission:

06 December 2019;

Accepted:

23 December 2019

doi: 10.17392/1101-20

Med Glas (Zenica) 2020; 17(1):106-109

INTRODUCTION

Rheumatic heart disease (RHD) is still the most common acquired heart disease worldwide. It causes permanent cardiac valve damage and leads to more severe conditions such as congestive heart disease, stroke, endocarditis, and death (1,2). Mortality rate from RHD is approximately 3.3%, annually. In Asia, RHD kills 356,000 – 542,000 patients every year (3,4). Rheumatic heart disease is a complex disease which is influenced by genetic factor, microorganism's virulence, and environmental factor (5).

An epidemiological study from 2009 found a relationship between genetic factor and RHD (6). Some human leukocyte antigen alleles are related to RHD (6). HLA-DR7 was reported to be a risk factor of RHD (2). A recent study in Uganda showed HLA alleles distribution in patients with RHD (7). The most common allele was HLA-DR11, which was found in 31.3% patients (7). Only 10.7% of healthy subjects in the study possessed HLA-DR11 allele (7).

Study regarding genetic factor, particularly HLA-DR alleles, and its relationship with RHD in Asia is scarce. Our study's goal was to provide the first data of HLA-DRB1 allele distribution among children with RHD in Medan, Indonesia as additional information for further investigation about the role of genetic factor in RHD.

PATIENTS AND METHODS

Patients and study design

An observational study was conducted in the Department of Child Health, Haji Adam Malik Hospital Medan (Indonesia) from April to June 2017. Patients were enrolled using consecutive sampling method. Inclusion criteria were children aged 5-18 years diagnosed with RHD. Children with concomitant congenital heart disease based on echocardiography evaluation were excluded. Children or proxies were interviewed to obtain gender and age data.

Informed consent was obtained before conducting the procedure.

An approval of the Ethics Committee, School of Medicine, Universitas Sumatera Utara was obtained.

Methods

Blood samples were drawn and genetic analysis was conducted for each sample. HLA-DRB1 alleles were analyzed using the PCR sequence-specific priming (SSP) technique using Olerup SSP® HLA typing kit (CareDx, Inc., Brisbane, California). Sample handling was conducted in Multidiscipline Laboratory Universitas Sumatera Utara.

Statistical analysis

Statistical analysis was done using computer software. Numerical data were presented in mean and standard deviation, whilst categorical ones in frequency and percentages.

RESULTS

A total of 62 children were enrolled in this study. Mean age of the children was 12.6 (SD 3.44) years with age range of 9-16. Males were more dominant, 33 (53.2%).

The most common allele was allele 12 (44 children), followed by allele 15 (32 children). The rarest allele was allele 09 (one child), followed by allele 04 (two children) (Figure 1).




Figure 1. HLA-DRB1 alleles distribution in 62 children with rheumatic heart disease

DISCUSSION

The RHD is common in developing countries and increases their national economic and epidemiologic burdens (2). In the next 5 to 20 years, approximately 15 million children and adolescents will suffer from RHD, and most of them require recurrent hospital admission and, not uncommonly, cardiothoracic surgery (8). The RHD is most prevalent in population between 5 and 16 years, hampering the crucial developmental period of a generation (9). As observed in this study, the mean age of RHD sufferer was 12.6 years. No

gender predilection is reported in the occurrence of RHD (10), but in our study male patients were slightly more dominant.

HLA plays an essential role in shaping one's immune system by regulating the antigen presentation of peptides and immune response (11). Therefore, it is suspected to be involved in autoimmune disease pathogenesis. Some autoimmune diseases such as Behcet's disease, celiac disease, and rheumatoid arthritis are related to HLA (11,12). The gene which codes a particular HLA protein in humans has several epitopes (12). The shared epitope for the HLA is passed on from generation to generation in an exclusive population. This is the reason why several populations are prone to a specific autoimmune disease, while some other are not. This proposed mechanism is called epitope hypothesis (13).

Similar condition is proposed in RHD. Host's genetic susceptibility has been suspected as one of the culprit of RHD (14,15). RHD is caused by immune response toward the Group A *Streptococcus* (GAS) infection (14). The microorganism often infects upper respiratory tract or skin. Several strains of GAS have surface proteins, especially protein M, which have similar molecular structure with cardiomyosin, tropomyosin, keratin, laminin, vimentin, and several cardiac valve proteins (10). Protein M is processed by HLA and presented to immune cells. Immune reaction is then triggered to eliminate the microorganism. Unfortunately, cross reaction between protein M and body's proteins also induces inflammation and damage in cardiac tissues (16). This pathogenesis confirms the concept of autoimmunity as the basis of RHD (2,17).

A study in India from 2010 to 2011 showed that several types of HLA were associated with RHD and the HLA type was different with other region's report (18). It can be inferred that HLA allele distribution in RHD patients is different

in each ethnicity. HLA-DR7 allele is the most common associated genetic predisposition for RHD. The allele is highly found in patients with RHD from Brazil-Mullatos and Brazil-Caucasians ethnicities (19). A study from India in 1986 showed a positive association between HLA-DR2 allele and RHD. On the other hand, HLA-DR2 allele was negatively associated with the disease (20). In Turkey, the presence of HLA-DR11 and HLA-A10 increased the risk of RHD (21). In Uganda, it was HLA-DR11 allele which increased the risk of RHD (8).

This is the first study to determine HLA-DRB1 allele distribution among children and adolescents with RHD. In our study, the most dominant allele is HLA-DRB1*12, followed by HLA-DRB1*15. Those findings are different from other studies from several regions and ethnicities. This is possibly caused by minor rearrangement in gene sequence at first. The rearrangement is induced by epigenetic events and accumulates as time goes by. This process creates different phenotypes from the ancient gene. The new gene sequence is then inherited and carried by next generation of specific populations (22).

A population based study is mandatory to extrapolate the result of this study, as this is a pilot hospital based study. Subjects from other regions of Indonesia should be involved to achieve more representative results.

In conclusion, it is proven in this research that RHD is influenced by genetic factor with HLA-DRB1*12 allele found to be the most common allele in children with RHD in Medan, Indonesia.

FUNDING

No specific funding was received for this study.

TRANSPARENCY DECLARATION

Conflict of interest: None to declare.

REFERENCES

1. Park, MK. Paediatric Cardiology for Practitioners: Acute Rheumatic Fever. 5th Ed., Philadelphia: Mosby Elsevier, 2008; 153-55.
2. Seckeler MD, Hoke TR. The worldwide epidemiology of acute rheumatic fever and rheumatic heart disease. Clin Epidemiol 2011; 3:67-84.
3. Carapetis JR. Rheumatic heart disease in Asia. Circulation 2008; 118:2748-53.
4. Kumar R, Raizada A, Aggarwal AK, Ganguly NK. A community-based rheumatic fever/ rheumatic heart disease cohort: twelve-year experience. Indian Heart J 2002; 54:54-8.
5. Beg A, Younas M, Asma T. Rheumatic heart disease: Socio-economic and environmental risk factors for acute rheumatic fever and rheumatic heart disease patients in Pakistan. Professional Med J 2016; 23:324-27.

6. Bryant PA, Robins-Browne R, Carapetis JR, Curtis N. Some of the people, some of the time. *Circulation* 2009; 119:742-53.
7. Okello E, Beaton A, Mondo KC, Kruzka P, Kiwanuka N, Adome-Odoi R, Freers J. Rheumatic heart disease in Uganda: the association between MHC classII HLA DR alleles and disease: a case control study. *BMC Cardiovasc Disord* 2014; 1-5.
8. Venugopal P, Gupta PK. Identification and diagnosis of risk factors and symptoms for rheumatic heart disease. *J Cardiovasc Dis Res* 2018; 9:109-14.
9. Prajapati D, Sharma D, Regmi PR, Khanal H, Baidya SG, Rajbhandari S, Rokka M, Singh S, Shrestha A, Shrestha A, Belbase, R, Sulpya, S. Epidemiological survey of rheumatic fever, rheumatic heart disease and congenital heart disease among school children in Kathmandu valley of Nepal. *Nepalese Heart Journal* 2014; 10:1-5.
10. Anderson RH. *Paediatric Cardiology: Rheumatic Fever*. 3rd edition. Philadelphia: Churchill Livingstone, 2010; 222-31.
11. Cortez A, Pulit SL, Leo PJ, Pointon JJ, Robinson PC, Weisman MH, Ward M, Gensler LS, Zhou X, Garchon H-J, Chiochia G, Nossent J, Lie BA, Forre O, Tuomilehto J, Laiho K, Bradbury LA, Elewaut D, Burgos-Vargas R, Stebbings S, Appleton L, Farrah C, Lau J, Haroon N, Mulero J, Blanco FJ, Gonzales-Gay MA, Lopez-Larrea C, Bowness P, Gaffney K, Gaston H, Gladman DD, Rahman P, Maksymowych WP, Crusius JBA, van der Horst-Bruinsma IE, Hansen IM, Inman RD, Martin J, Breban M, Wordsworth BP, Reveille JD, Evans DM, Bkker PIW, Brown MA. Major histocompatibility complex associations of ankylosing spondylitis are complex and involve further epistasis with ERAP1. *Nat Commun* 2015; 6:7146.
12. Komata T, Tsuchiya N, Hatta Y, Matsushita M, Ogawa A, Mitsui H, Tokunaga K. HLA-DRB1 alleles encoding the shared epitope associated with rheumatoid arthritis confer additional susceptibility to seronegative spondyloarthropathies in HLA-B27-positive Japanese individuals. *Jpn J Rheumatol* 1999; 9:55-64.
13. Bodis G, Toth V, Schwarting A. Role of human leukocyte antigens (HLA) in autoimmune diseases. *Rheumatol Ther* 2018; 5:5-20.
14. Carapetis JR, Mc Donald M, Wilson NJ. Acute rheumatic fever. *Lancet* 2005; 336:155-68.
15. Steer AC, Carapetis JR, Nolan TM, Shann F. Systematic review of rheumatic heart disease prevalence in children in developing countries: the role of environmental factors. *Circulation* 2002; 38:229-34.
16. Guilherme L, Kalil J. Rheumatic fever and rheumatic heart disease: cellular mechanisms leading autoimmune reactivity and disease. *J Clin Immunol* 2010; 30:17-23.
17. Fae K, Oshiro SE, Toubert A, Charron D, Kalil J, Guilherme L. How an autoimmune reaction triggered by molecular mimicry between streptococcal M protein. *J Autoimmun* 2005; 24:101-9.
18. Bajoria D. The HLA class II associations with rheumatic heart disease in south Indian patients. *J Clin Diag Res* 2003; 7:302-4.
19. Guilherme L, Kalil J, Cunningham M. Molecular mimicry in the autoimmune pathogenesis of RHD. *Autoimmunity* 2006; 39:31-9.
20. Jhinghan B, Mehra K, Reddy KS, Taneja V, Vaidya MC, Bhatia ML. HLA, blood groups, and secretor status in patients with established rheumatic fever and rheumatic heart disease. *Tissue Antigens* 1986; 27:172-8.
21. Olmez U, Turgay M, Ozenirler S, Tutkak H, Duzgun N, Duman M, Tokgoz G. Association of HLA class I and class II antigens with rheumatic fever in a Turkish population. *Scand J Rheumatol* 1993; 22:49-52.
22. Guilherme L, Kohler KF, Postol E, Kalil J. Genes, autoimmunity and pathogenesis of rheumatic heart disease. *Ann Pediatr Cardiol* 2011; 4:13-21.